

D-Branes and Derived Categories

Yuri Malyuta

*Institute for Nuclear Research
National Academy of Sciences of Ukraine
03022 Kiev, Ukraine
e-mail: interdep@kinr.kiev.ua*

Abstract

The digest of ideology interpreting D-branes on Calabi-Yau manifolds as objects of the derived category is given.

Keywords: D-branes, Derived category, Triangulated structure,
Monodromy.

1 Introduction

Recently there has been substantial progress [1, 2, 3] in understanding D-branes on Calabi-Yau manifolds in context of derived categories [4].

The purpose of the present paper is to give the digest of this ideology.

2 Sheaves

In this section we shall introduce the definitions of presheaves and sheaves [5].

A *presheaf* F over a topological space X is

- 1) An assignment to each nonempty open set $U \subset X$ of a set $F(U)$ (*sections* of a presheaf F);
- 2) A collection of mappings (called restriction homomorphisms)

$$r_{UV} : F(U) \rightarrow F(V)$$

for each pair of open sets U and V such that $V \subset U$, satisfying

$$r_{UU} = 1_U, \quad r_{VW} r_{UV} = r_{UW} \quad \text{for } W \subset V \subset U.$$

A presheaf F is called a *sheaf* if for every collection U_i of open subsets of X with $U = \bigcup_{i \in I} U_i$ the following axioms hold :

- a) If $s, t \in F(U)$ and $r_{UU_i}(s) = r_{UU_i}(t)$ for all i , then $s = t$;
- b) If $s_i \in F(U_i)$ and if for $U_i \cap U_j \neq \emptyset$ we have

$$r_{U_i, U_i \cap U_j}(s_i) = r_{U_j, U_i \cap U_j}(s_j)$$

for all i , then there exists an $s \in F(U)$ such that $r_{U,U_i}(s) = s_i$ for all i .

If F and G are presheaves over X , then a *morphism* of presheaves $f : F \rightarrow G$ is a collection of maps $f(U) : F(U) \rightarrow G(U)$, satisfying the relation $r_{UV} f(U) = f(V) r_{UV}$.

Morphisms of sheaves are simply morphisms of the underlying presheaves.

Let $(X, \mathcal{O})$ be a complex manifold. A sheaf B over X is called a *coherent sheaf* of $\mathcal{O}$ -modules if for each $x \in X$ there is a neighborhood U of x such that there is an exact sequence of sheaves over U ,

$$0 \rightarrow B|_U \rightarrow \mathcal{O}^{\oplus p_1}|_U \rightarrow \mathcal{O}^{\oplus p_2}|_U \rightarrow \dots \rightarrow \mathcal{O}^{\oplus p_k}|_U \rightarrow 0.$$

3 Complexes

Let $B^\bullet$ denote a *complex* of coherent sheaves [4]

$$B^\bullet : \dots \xrightarrow{d^{i-2}} B^{i-1} \xrightarrow{d^{i-1}} B^i \xrightarrow{d^i} B^{i+1} \xrightarrow{d^{i+1}} \dots,$$

where $d^i d^{i-1} = 0$.

Cohomology groups of the complex $B^\bullet$ are defined as

$$H^i(B^\bullet) = \text{Ker } d^i / \text{Im } d^{i-1}.$$

A morphism of complexes $f : B^\bullet \rightarrow C^\bullet$ induces a morphism of cohomology groups $H(f) : H^\bullet(B^\bullet) \rightarrow H^\bullet(C^\bullet)$.

If $H(f)$ is an isomorphism, the morphism f is said to be a *quasi-isomorphism*.

If morphisms f and g are *homotopy equivalent*, then $H(f) = H(g)$.

4 Categories

In this section we shall give some formal definitions [6].

A *category* $\mathcal{C}$ consists of the following data :

- 1) A class $\text{Ob } \mathcal{C}$ of objects $A, B, C, \dots$;
- 2) A family of disjoint sets of morphisms $\text{Hom}(A, B)$, one for each ordered pair A, B of objects ;
- 3) A family of maps

$$\text{Hom}(A, B) \times \text{Hom}(B, C) \rightarrow \text{Hom}(A, C) ,$$

one for each ordered triplet A, B, C of objects.

These data obey the axioms :

- a) If $f : A \rightarrow B$, $g : B \rightarrow C$, $h : C \rightarrow D$, then composition of morphisms is associative, that is, $h(gf) = (hg)f$;
- b) To each object B there exists a morphism $1_B : B \rightarrow B$ such that $1_B f = f$, $g 1_B = g$ for $f : A \rightarrow B$ and $g : B \rightarrow C$.

An *additive category* is a category in which each set of morphisms $\text{Hom}(A, B)$ has the structure of an abelian group, subject to the following axioms :

A1 Composition of morphisms is distributive, that is,

$$(g_1 + g_2)f = g_1f + g_2f , \quad h(g_1 + g_2) = hg_1 + hg_2$$

for any $g_1, g_2 : B \rightarrow C$, $f : A \rightarrow B$, $h : C \rightarrow D$;

A2 There is a null object 0 such that $\text{Hom}(A, 0)$ and $\text{Hom}(0, A)$ consist of one morphism for any A ;

A3 To each pair of objects A_1 and A_2 there exists an object B and four morphisms

$$\begin{array}{ccccc} & p_1 & & p_2 & \\ & \swarrow & & \searrow & \\ A_1 & \xleftrightarrow{\quad} & B & \xleftrightarrow{\quad} & A_2 \\ & \nwarrow & & \nearrow & \\ & i_1 & & i_2 & \end{array}$$

which satisfy the identities

$$p_1 i_1 = 1_{A_1} , \quad p_2 i_2 = 1_{A_2} , \quad i_1 p_1 + i_2 p_2 = 1_B , \quad p_2 i_1 = p_1 i_2 = 0 .$$

An *abelian category* $\mathcal{A}$ is an additive category which satisfies the additional axiom :

A4 To each morphism $f : A \rightarrow B$ there exists the sequence

$$K \xrightarrow{k} A \xrightarrow{i} I \xrightarrow{j} B \xrightarrow{c} K'$$

with the properties

- a) $ji = f$,
- b) K is a kernel of f , K' is a cokernel of f ,
- c) I is a cokernel of k and a kernel of c .

The *category of coherent sheaves* is the abelian category $\mathcal{A}$.

5 The derived category

The derived category $D(\mathcal{A})$ is constructed in three steps [4] :

1st step. The *category of complexes* of coherent sheaves $\text{Kom}(\mathcal{A})$ is determined as follows

$$\text{Ob } \text{Kom}(\mathcal{A}) = \{\text{complexes } B^\bullet \text{ of coherent sheaves}\} ,$$

$$\text{Hom}(B^\bullet, C^\bullet) = \{\text{morphisms of complexes } B^\bullet \rightarrow C^\bullet\} ;$$

2nd step. The *homotopy category* $K(\mathcal{A})$ is determined as follows

$$\text{Ob } K(\mathcal{A}) = \text{Ob } \text{Kom}(\mathcal{A}) ,$$

$$\text{Mor } K(\mathcal{A}) = \text{Mor } \text{Kom}(\mathcal{A}) \text{ modulo homotopy equivalence ;}$$

3rd step. The *derived category* $D(\mathcal{A})$ is determined as follows

$$\text{Ob } D(\mathcal{A}) = \text{Ob } K(\mathcal{A}) ,$$

The morphisms of $D(\mathcal{A})$ are obtained from morphisms in $K(\mathcal{A})$ by inverting all quasi-isomorphisms.

The derived category $D(\mathcal{A})$ is the additive category.

6 Triangulated structure


The derived category $D(\mathcal{A})$ admits a *triangulated structure* [4] with *shift functor* $[n]$ defined by

$$(B[n])^i = B^{n+i}$$

and with a class of *distinguished triangles*

$$\begin{array}{ccc} & C & \\ \swarrow^{[1]} & & \nwarrow \\ A & \longrightarrow & B \end{array} \quad C = A[1] \oplus B$$

These data satisfy a number of axioms. The *octahedral axiom* is an essential ingredient in the study of D-brane stability [1]. The octahedral axiom states that there exists the octahedron consisting of a top cap and a bottom cap :



(distinguished triangles are marked by $\bullet$)

Interpreting D-branes as vertices of the octahedron, it is possible to describe *D-brane decays* : if C is stable against decay into A and B , but that B itself is unstable with respect to a decay into E and F , then C will always be unstable with respect to decay into F and some bound state G of A and E .

7 The quintic

Let X be the quintic hypersurface in CP^4 . The mirror Y is defined as the orbifold $X/\mathbf{Z}_5^3$. In virtue of *mirror symmetry* [7] the *Kähler moduli space* of X is identified with the *complex structure moduli space* of Y . The complex structure moduli space of Y is described by the Picard-Fuchs equation

$$\{\theta_z^4 + 5z(5\theta_z + 4)(5\theta_z + 3)(5\theta_z + 2)(5\theta_z + 1)\}\omega_k(z) = 0 ,$$

where $\theta_z = z d/dz$, the complex variable z spans the complex structure moduli space of Y .

The *Landau-Ginzburg point* of the moduli space of X is mirror to $z = \infty$, the *large radius limit* of X is mirror to $z = 0$, the *conifold point* of X is mirror to $z = 1$. The periods $\omega_k(z)$ are singular at these three points.

8 Monodromy

Acting on the derived category $D(\mathcal{A})$, the *monodromy* is induced by a *Fourier-Mukai transform* [8] associated to some generator $K^\bullet \in D(\mathcal{A})$. The formula for the monodromy action on a complex $B^\bullet$ is

$$B^\bullet \longmapsto Rp_{1*}(K^\bullet \overset{L}{\otimes} p_2^*(B^\bullet)) .$$

Geometry associated to this monodromy action is

$$\begin{array}{ccc} & \triangle & \\ & \cap & \\ & X \times X & \\ p_1 \swarrow & & \searrow p_2 \\ X & & X \end{array}$$

where $\triangle \subset X \times X$ is the diagonal embedding of X .

In the formula for the monodromy action, we

- 1) Take a complex of sheaves $B^\bullet$ on X , "pull it back" to the inverse-image complex of sheaves $p_2^*(B^\bullet)$ on $X \times X$;
- 2) Take the tensor-product with the generator $K^\bullet$ and construct the left-derived complex of sheaves ;
- 3) "Push-forward" to the direct image complex $p_{1*}(\cdot)$ and construct the right-derived complex of sheaves on X .

The most obvious monodromy is that about the Landau-Ginsburg point in the Kähler moduli space of the quintic. This monodromy is generated by [1]

$$K_{LG}^\bullet = 0 \rightarrow \mathcal{O} \boxtimes \mathcal{O}(1) \rightarrow \mathcal{O}_\Delta(1) \rightarrow 0.$$

The monodromy calculations for $\mathcal{O} \in D(\mathcal{A})$ yield the result

$$\begin{aligned} M_{LG}(\mathcal{O}) &= 0 \rightarrow \mathcal{O}^{\oplus 5} \rightarrow \mathcal{O}(1) \rightarrow 0 \\ (M_{LG})^2(\mathcal{O}) &= 0 \rightarrow \mathcal{O}^{\oplus 10} \rightarrow \mathcal{O}(1)^{\oplus 5} \rightarrow \mathcal{O}(2) \rightarrow 0 \\ (M_{LG})^3(\mathcal{O}) &= 0 \rightarrow \mathcal{O}^{\oplus 10} \rightarrow \mathcal{O}(1)^{\oplus 10} \rightarrow \mathcal{O}(2)^{\oplus 5} \rightarrow \mathcal{O}(3) \rightarrow 0 \\ (M_{LG})^4(\mathcal{O}) &= \mathcal{O}(-1)[4] \\ (M_{LG})^5(\mathcal{O}) &= \mathcal{O}[2] \end{aligned}$$

9 Boundary linear σ -model

Boundary linear σ -model [2] is determined by the Lagrangian

$$\begin{aligned} L = \sum_n & \left(i\bar{\beta}^{(2n)} \partial_0 \beta^{(2n)} + i\bar{\rho}^{(2n+1)} \partial_0 \rho^{(2n+1)} + \right. \\ & + \frac{1}{2} \bar{\beta}^{(2n)} (|\kappa^{(2n+1)}|^2 \bar{c}_{2n} c_{2n} + |\kappa^{(2n)}|^2 c_{2n-1} \bar{c}_{2n-1}) \beta^{(2n)} + \\ & \left. + \frac{1}{2} \bar{\rho}^{(2n+1)} (|\kappa^{(2n+2)}|^2 \bar{c}_{2n+1} c_{2n+1} + |\kappa^{(2n+1)}|^2 c_{2n} \bar{c}_{2n}) \rho^{(2n+1)} \right), \end{aligned}$$

which involves superfields $\beta^{(2n)}$, $\rho^{(2n+1)}$, c_k .

Consider the complex of direct sums of holomorphic line bundles

$$\dots \rightarrow \bigoplus_i \mathcal{O}(m_i^{(2n-1)}) \xrightarrow{c_{2n-1}} \bigoplus_i \mathcal{O}(m_i^{(2n)}) \xrightarrow{c_{2n}} \bigoplus_i \mathcal{O}(m_i^{(2n+1)}) \xrightarrow{c_{2n+1}} \dots$$

Sections of holomorphic line bundles describe superfields $\beta^{(2n)}$, $\rho^{(2n+1)}$; differentials describe superfields c_k .

Acknowledgement

This material was presented in the lecture given by the author at the Institute of Mathematics (Kiev, Ukraine). The author thanks audience for questions and comments.

References

- [1] P.S. Aspinwall and M.R. Douglas, *D-brane stability and monodromy*, hep-th/0110071.
- [2] J. Distler, H. Jockers and H. Park, *D-brane monodromies, derived categories and boundary linear sigma models*, hep-th/0206242.
- [3] P.S. Aspinwall, R.L. Karp and R.P. Horja, *Massless D-branes on Calabi-Yau threefolds and monodromy*, hep-th/0209161.
- [4] S.I. Gelfand and Yu.I. Manin, *Homological algebra*, Springer-Verlag, Berlin, 1994.
- [5] R.O. Wells, *Differential analysis on complex manifolds*, Springer-Verlag, Berlin, 1980.
- [6] S. MacLane, *Categorical algebra*, Bull. Amer. Math. Soc. **71** (1965) 40.
- [7] P. Candelas, X.C. de la Ossa, P.S. Green and L. Parkes, *A pair on Calabi-Yau manifolds as an exactly soluble superconformal theory*, Nucl. Phys. **B359** (1991) 21.
- [8] M.R. Douglas, *D-branes, categories and $N=1$ supersymmetry*, hep-th/0011017.